## STATUTORY RULES OF NORTHERN IRELAND

#### 2002 No. 217

#### **FOOD**

# Meat (Hazard Analysis and Critical Control Point) Regulations (Northern Ireland) 2002

*Made* . . . . . . . 10th June 2002

Coming into operation in accordance with regulation 2

The Department of Health, Social Services and Public Safety(a), in exercise of the powers conferred on it by Articles 15(1)(b) and (f) and 16(1) of the Food Safety (Northern Ireland) Order 1991(b), and being a Department designated(c) for the purposes of section 2(2) of the European Communities Act 1972(d) in relation to the common agricultural policy of the European Community, in exercise of the powers conferred on it by the said section 2(2), and having had regard in accordance with Article 47(3A) of the said Order to relevant advice given by the Food Standards Agency (in so far as the Regulations are made in exercise of the powers conferred by the said provisions of the said Order) and after consultation in accordance with Article 47(3) and (3B) of the said Order with such organisations as appear to it to be representative of interests likely to be substantially affected by the Regulations, and after consultation as required by Article 9 of Regulation (EC) No. 178/2002 of the European Parliament and of the Council laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety(e), and in exercise of all other powers enabling it in that behalf, hereby makes the following Regulations:

# Citation, interpretation and scope

- **1.**—(1) These Regulations may be cited as the Meat (Hazard Analysis and Critical Control Point) Regulations (Northern Ireland) 2002.
  - (2) In these Regulations—
  - "the Fresh Meat Regulations" means the Fresh Meat (Hygiene and Inspection) Regulations (Northern Ireland) 1997(**f**);
  - "fresh poultry meat" has the same meaning as in regulation 2(1) of the Poultry Meat Regulations;
  - "fresh red meat" means fresh meat as defined in regulation 2(1) of the Fresh Meat Regulations;
  - "licensed poultry meat cutting plant" means cutting premises licensed under regulation 4 of the Poultry Meat Regulations;
  - "licensed poultry meat slaughterhouse" means a slaughterhouse licensed under regulation 4 of the Poultry Meat Regulations;
  - "licensed red meat cutting plant" means cutting premises licensed under regulation 4 of the Fresh Meat Regulations;
  - "licensed red meat slaughterhouse" means a slaughterhouse licensed under regulation 4 of the Fresh Meat Regulations;
  - "livestock unit" has the same meaning as in regulation 2(1) of the Fresh Meat Regulations;
  - "poultry" means domestic fowls, turkeys, guinea fowls, ducks and geese;

(a) Formerly the Department of Health and Social Services; see S.I. 1999/283 (N.I. 1), Article 3

<sup>(</sup>b) S.I. 1991/762 (N.I. 7) as amended by S.I. 1996/1633 (N.I. 12) and paragraphs 26 to 42 of Schedule 5 and Schedule 6 to the Food Standards Act 1999 c. 28. Regulation 13(4) of S.R. 2000 No. 78 expressly authorises the Department of Health, Social Services and Public Safety to amend or revoke existing Regulations made or having effect as if made by the Department of Agriculture and Rural Development (including those made jointly with the Department of Health, Social Services and Public Safety) under the Food Safety (Northern Ireland) Order 1991

c) S.I. 2000/2812

<sup>(</sup>d) 1972 c. 68

<sup>(</sup>e) O.J. No. L31, 1.2.2002, p. 1

<sup>(</sup>f) S.R. 1997 No. 493; the relevant amending Regulations are S.R. 2000 Nos. 78 and 191

"the Poultry Meat Regulations" means the Poultry Meat, Farmed Game Bird Meat and Rabbit Meat (Hygiene and Inspection) Regulations (Northern Ireland) 1995(a); and "the specified amount" means—

- (a) in the case of any licensed red meat slaughterhouse, 500 livestock units;
- (b) in the case of any licensed red meat cutting plant, 150 tonnes of fresh red meat;
- (c) in the case of any licensed poultry meat slaughterhouse, 200,000 poultry; and
- (d) in the case of any licensed poultry meat cutting plant, 150 tonnes of fresh poultry meat.
- (3) Nothing in these Regulations shall apply in relation to any licensed red meat slaughterhouse which is used to process bovine animals under the purchase scheme introduced by Commission Regulation (EC) No. 716/96(b) adopting exceptional support measures for the beef market in the United Kingdom.

#### Commencement

- **2.**—(1) Subject to paragraph (2), these Regulations shall come into operation on 15th July 2002.
- (2) Notwithstanding paragraph (1), these Regulations shall come into operation on 7th June 2003 in so far as they apply in relation to any small meat establishment.
  - (3) For the purposes of paragraph (2), "small meat establishment" means—
  - (a) a licensed red meat slaughterhouse, a licensed poultry meat slaughterhouse, a licensed red meat cutting plant or a licensed poultry meat cutting plant which—
    - (i) was operational in the calendar year 2000 and had an average weekly throughput in that year lower than the specified amount,
    - (ii) became operational for the first time on or after 1st January 2001 but before 1st May 2002 and had an average weekly throughput during the time it was operational in that period lower than the specified amount, or
    - (iii) became operational for the first time on or after 1st May 2002 and which the Food Standards Agency considers is likely to have an average weekly throughput lower than the specified amount;
  - (b) a cold store licensed under regulation 4 of the Fresh Meat Regulations or regulation 4 of the Poultry Meat Regulations, in each case with a storage capacity of less than 25,000 cubic metres;
  - (c) a re-packaging centre licensed under regulation 4 of the Fresh Meat Regulations; or
  - (d) a re-wrapping centre licensed under regulation 4 of the Poultry Meat Regulations.

# Amendments to the Fresh Meat (Hygiene and Inspection) Regulations (Northern Ireland) 1997

- **3.**—(1) The Fresh Meat Regulations are amended in accordance with the following paragraphs.
- (2) The following sub-paragraph is inserted between sub-paragraphs (aA) and (b) of paragraph (1) of regulation 7 (supervision of premises)—
  - "(aB) the inspection of any documents and records required to be retained by the occupier pursuant to regulation 19(1)(eA) or (eB);".
- (3) In paragraph (1)(e) of regulation 7, for "and 17" there shall be substituted ", 17, 17A, 17B and 17C".
  - (4) In sub-paragraph (d) of paragraph (1) of regulation 19 (duties of occupier)—
  - (a) for "carry out" there shall be substituted "conduct regular"; and
  - (b) the phrase "(including any microbiological checks the Agency may require)" is revoked.
- (5) In sub-paragraph (e) of paragraph (1) of regulation 19, for "pursuant to" there shall be substituted "by virtue of".

<sup>(</sup>a) S.R. 1995 No. 396, as amended by S.R. 1997 No. 496, S.R. 1998 No. 237, S.R. 2000 Nos. 78 and 191 and S.R. 2001 No. 429

<sup>(</sup>b) O.J. No. L99, 20.4.96, p. 14

- (6) The following sub-paragraphs are inserted between sub-paragraphs (e) and (f) of paragraph (1) of regulation 19—
  - "(eA) shall retain for a period of at least one year any documents and records established by him in accordance with paragraph (4)(g);
  - (eB) shall retain for a period of at least 18 months any records which, in compliance with Schedule 17A, 17B or 17C, as appropriate, he has made following the carrying out by him of microbiological checks in accordance with paragraph (5);".
  - (7) The following paragraphs are added after paragraph (3) of regulation 19—
  - "(4) The occupier of any licensed slaughterhouse, licensed cutting premises, licensed cold store or licensed repackaging centre shall conduct the regular checks on the general hygiene of conditions of production in those premises which are required by paragraph (1)(d) by implementing and maintaining a permanent procedure developed in accordance with the following principles—
    - (a) identify any hazards that must be prevented, eliminated or reduced to acceptable levels;
    - (b) identify the critical control points at the step or steps at which control is essential to prevent or eliminate a hazard or reduce it to acceptable levels;
    - (c) establish critical limits at critical control points which separate acceptability from unacceptability for the prevention, elimination or reduction of identified hazards;
    - (d) establish and implement effective monitoring procedures at critical control points;
    - (e) establish corrective actions when monitoring indicates that a critical control point is not under control;
    - (f) establish procedures to verify whether the measures outlined in sub-paragraphs
       (a) to (e) are working effectively; verification procedures shall be carried out regularly; and
    - (g) establish documents and records commensurate to the nature and size of the business to demonstrate the effective application of the measures outlined in sub-paragraphs (a) to (f) and to facilitate official controls.
  - (5) The occupier of any licensed slaughterhouse shall, in conducting the regular checks on the general hygiene of conditions of production in those premises which are required by paragraph (1)(d), carry out microbiological checks—
    - (a) in relation to carcases, in accordance with the procedures laid down in—
      - (i) Schedule 17A, or
      - (ii) Schedule 17B; and
    - (b) in relation to cleaning and disinfection of the premises, in accordance with the procedures laid down in Schedule 17C.
  - (6) The occupier of any licensed cutting premises shall, in conducting the regular checks on the general hygiene of conditions of production in those premises which are required by paragraph (1)(d), carry out microbiological checks in relation to cleaning and disinfection of the premises, in accordance with the procedures laid down in Schedule 17C.".
  - (8) The following Schedules are inserted between Schedules 17 and 18—

"SCHEDULE 17A

Regulations 7(1)(e), 19(1)(eB) and 19(5)(a)(i)

#### Community procedures for conducting microbiological checks on carcases

#### Sampling procedure and number of samples to be taken

1. (a) Between 5 and 10 carcases should be sampled on a single day during each week. The day of sampling should be changed to ensure that every day of the working week is covered. The frequency of sampling the carcases in low-throughput slaughterhouses and in slaughterhouses not working on a full-time basis should

- be determined by the OVS based on his judgement on hygiene standards with respect to the slaughter at each plant.
- (b) A sample from four sites from each carcase should be taken half way through the slaughter day, after dressing and before chilling.
- (c) Carcase identification, date and time of sampling should be recorded for each sample and the name of the person performing the sampling.
- (d) The frequency of sampling may be reduced to fortnightly testing if satisfactory results are obtained on six consecutive weeks, but weekly sampling must be resumed if unsatisfactory results are obtained.

## Sampling sites

2. (a) The following sites will usually be appropriate for process control:

Cattle: neck, brisket, flank, and rump

Sheep, goat: flank, thorax lateral, brisket, and breast

Pig: back, jowl (or cheek), hind limb medial (ham), and belly

Horse: flank, brisket, back, and rump.

(b) However, alternative sites may be used, following consultation with the OVS where it has been demonstrated that, because of the slaughter technology at a particular plant, other sites are more likely to carry higher levels of contamination. In these cases, sites shown to carry higher levels of contamination may be chosen.

## **Excision sampling method**

- 3. The following protocol should be followed at the slaughterhouse:
  - (a) Four tissue samples representing a total of 20 cm<sup>2</sup> should be obtained from each carcase.
  - (b) Pieces of tissue may be obtained using a sterile cork borer (2.5 cm diameter) or by cutting a slice of 5 cm<sup>2</sup> and maximum thickness of 5 mm off the carcase with a sterile instrument.
  - (c) Samples from the four sampling sites of each tested carcase may be analysed separately or may be pooled in the same container before examination. Where unacceptable results are obtained from pooled samples and corrective actions do not lead to better hygiene, further samples should not be pooled until problems have been resolved.
  - (d) The samples must be placed aseptically into a sample container or plastic dilution bag at the slaughterhouse, for transfer to the laboratory.

#### Method for the examination of samples

- 4. The following protocol should be followed in the laboratory:
  - (a) Samples should be stored refrigerated until examination at 4°C. Samples should be examined within 24 hours after sampling.
  - (b) Samples should be homogenised in a plastic dilution bag for at least two minutes in 100 ml of dilution media (see ISO 6887–1) at about 250 cycles of a peristaltic Stomacher or homogenised by a rotary blender (homogeniser).
  - (c) Dilution before plating should be carried out in 10-fold steps in the dilution media.
  - (d) Analysis should be performed for total viable counts and Enterobacteriaceae. ISO methods should provide the basis for examination of samples.

#### Records

- 5. (a) All test results must be recorded in terms of colony forming units (cfu) per cm<sup>2</sup> of surface area. The daily log mean results for carcases sampled on one day must be calculated and recorded.
  - (b) Records must include:
    - (i) type, origin and identification of the sample, date and time of sampling, name of the person that performed the sampling,

- (ii) name and address of the laboratory which analysed the sample, date of investigation of samples in the laboratory and details of the method used including inoculation of different agars, incubation temperature, time, and results as number of cfu per plate used to calculate the result in cfu/cm<sup>2</sup> of surface area.
- (c) A responsible person from the laboratory should sign the records.
- (d) To permit evaluation, results must be shown on process control charts or tables, containing at least the last 13 weekly test results in order.

#### Verification criteria

- 6. (a) Daily log mean results must be allocated into one of three categories for process control verification: "acceptable", "marginal", and "unacceptable" as set out in the table, where 'M' and 'm' denote the upper limits for the marginal and acceptable categories, respectively, for samples taken according to the excision method.
  - (b) The test results should be categorised according to the respective microbiological criteria in the same order as the samples are collected.
  - (c) As each new test result is obtained, the verification criteria are applied anew to evaluate the status of process control with respect to microbiological contamination and hygiene.
  - (d) An unacceptable result or unsatisfactory marginal result trends should trigger action to review process controls, discover the cause if possible, and prevent recurrence.

Daily log mean values (cfu/cm <sup>2</sup> )	Acceptable range		Marginal range (>m but <m)< th=""><th>Unacceptable range (&gt;M)</th></m)<>	Unacceptable range (>M)
	Cattle/sheep/ goat/horse	Pig	Cattle/pig/sheep/goat/ horse	Cattle/pig/sheep goat/horse
Total viable counts (TVC)	<3.5 log	<4.0 log	<3.5 log (pig: 4.0 log) - 5.0 log	>5.0 log
Enterobacteriaceae	<1.5 log	<2.0 log	1.5 log (pig: 2.0 log) - 2.5 log (pig: 3.0 log)	>2.5 log (pig >3.0 log)

#### Feedback to staff

- 7. (a) The results of the test must be fed back to the responsible staff as soon as possible.
  - (b) The results should be used to maintain and improve the standard of slaughter hygiene. Causes of poor results may be clarified by consultation with the slaughtering staff where the following factors could be involved: poor working procedures, absence or inadequacy of training and/or instructions, the use of unsuitable cleaning and/or disinfection materials and chemicals, inadequate maintenance of cleaning apparatus, and inadequate supervision.

SCHEDULE 17B

Regulations 7(1)(e), 19(1)(eB) and 19(5)(a)(ii)

## National procedures for conducting microbiological checks on carcases

### Sampling procedure and number of samples to be taken

1. (a) Between 5 and 10 carcases should be sampled on a single day each week. The day of sampling should be changed each week to ensure that every day of the working week is covered. The frequency of testing the carcases in low-throughput slaughterhouses and for slaughterhouses not working on a full-time basis should be determined by the OVS based on his judgement on hygiene standards with respect to the slaughter at each plant.

- (b) A sample from four sites from each carcase should be taken half way through the slaughter day, after dressing and before chilling.
- (c) Carcase identification, date and time of sampling should be recorded for each sample and the name of the person performing the sampling.
- (d) The frequency of sampling may be reduced to fortnightly testing if satisfactory results are obtained on six consecutive weeks, but weekly sampling must be resumed if unsatisfactory results are obtained.

# Sampling sites

2. (a) The following sites will usually be appropriate for process control:

Cattle: neck, brisket, flank, and rump

Sheep, goat: flank, thorax lateral, brisket, and breast

Pig: back, jowl (or cheek), hind limb medial (ham), and belly

Horse: flank, brisket, back, and rump.

(b) However, alternative sites may be used, following consultation with the OVS where it has been demonstrated that, because of the slaughter technology at a particular plant, other sites are more likely to carry higher levels of contamination. In these cases sites shown to carry higher levels of contamination may be chosen.

# Wet and dry swabbing method

- 3. The following protocol should be followed at the slaughterhouse:
  - (a) Where swabs are moistened prior to collection of samples, a sterile peptone salt diluent (see ISO 6887–1) should be used.
  - (b) The sampling area for swabbing should cover 100 cm<sup>2</sup> for cattle and horses, 50 cm<sup>2</sup> for pigs, sheep and goats per sampling site. However, a smaller area may be tested, subject to the approval of the OVS on the basis of historical data, and a minimum area of 10 cm<sup>2</sup> per site.
  - (c) The swab should be moistened for at least 5 seconds in the diluent and rubbed initially vertically, then horizontally, then diagonally for not less than 20 seconds across the swab site. As much pressure as possible should be used. Repeat swabbing method using a dry swab at the same site.
  - (d) Samples from the four sampling sites of each tested carcase may be analysed separately or may be pooled in the same container before examination. Where unacceptable results are obtained with pooled samples and corrective actions do not lead to better hygiene, further samples should not be pooled until problems have been resolved.
  - (e) The samples must be placed aseptically into a sample container or plastic dilution bag at the slaughterhouse for transfer to the laboratory.

### Method for the examination of samples

- 4. The following protocol should be followed in the laboratory:
  - (a) Samples should be stored refrigerated until examination at 4°C. Samples should be examined within 24 hours after sampling.
  - (b) Samples should be homogenised in a plastic dilution bag for at least two minutes in 100 ml of dilution media (see ISO 6887–1) at about 250 cycles of a peristaltic Stomacher or homogenised by a rotary blender (homogeniser). Alternatively swab samples may be shaken vigorously in the dilution media.
  - (c) Dilution before plating should be carried out in 10-fold steps in the dilution media.
  - (d) Analysis should be performed for total viable counts and Enterobacteriaceae. ISO methods should provide the basis for examination of samples.

#### Records

5. (a) All test results must be recorded in terms of colony forming units (cfu) per cm<sup>2</sup> of surface area. The daily log mean results for the carcases sampled on one day must be calculated and recorded.

- (b) Records must include:
  - (i) Origin, type and identification of the sample, date and hour of sampling, name of the person that performed the sampling.
  - (ii) Name and address of the laboratory which analysed the sample, date of investigation of samples in the laboratory and details of the method used including inoculation of different agars, incubation temperature, time, and results as number of cfu per plate used to calculate the result in (cfu) per cm<sup>2</sup> of surface area.
- (c) A responsible person from the laboratory should sign the records.
- (d) To permit evaluation, results must be shown on process control charts or tables, containing at least the last 13 weekly test results in order.

#### Verification criteria

- 6. (a) Daily log mean results must be allocated into one of three categories for process control verification: "acceptable", "marginal", and "unacceptable" as set out in the table, where 'M' and 'm' denote the upper limits for the marginal and acceptable categories, respectively, for samples taken according to the wet and dry swabbing method.
  - (b) The test results should be categorised according to the respective microbiological criteria in the same order as the samples are collected.
  - (c) As each new test result is obtained, the verification criteria are applied anew to evaluate the status of process control with respect to microbiological contamination and hygiene.
  - (d) An unacceptable result or unsatisfactory marginal result trends should trigger action to review process controls, discover the cause if possible, and prevent recurrence.

Daily log mean values (cfu/cm <sup>2</sup> )	Acceptable range		Marginal range (>m but <m)< th=""><th>Unacceptable range (&gt;M)</th></m)<>	Unacceptable range (>M)
	Cattle/sheep/ goat/horse	Pig	Cattle/pig/sheep/goat/horse	Cattle/pig/sheep/goat/horse
Total viable counts (TVC)	<2.8 log	<3.3 log	2.8 log (pig: 3.3 log) - 4.3 log	>4.3 log
Enterobacteriaceae	<0.8 log	<1.3 log	0.8 log (pig: 1.3 log) -1.8 log (pig: 2.3 log)	>1.8 log (pig: >2.3 log)

#### Feedback to staff

- 7. (a) The results of the test must be fed back to the responsible staff as soon as possible.
  - (b) The results should be used to maintain and improve the standard of slaughter hygiene. Causes of poor results may be clarified by consultation with the slaughtering staff where the following factors could be involved: poor working procedures, absence or inadequacy of training and/or instructions, the use of unsuitable cleaning and/or disinfection materials and chemicals, inadequate maintenance of cleaning apparatus, and inadequate supervision.

SCHEDULE 17C

Regulations 7(1)(e), 19(1)(eB) and 19(5)(b) and (6)

# Community procedures for conducting microbiological checks in relation to cleaning and disinfection of premises

1. Microbiological sampling must take place before production starts, never during production. If visible dirt is present, cleaning should be judged as unacceptable without any further microbiological evaluation.

### Sampling sites

- 2. (a) To ensure that all surfaces are tested in the course of a month a schedule should be made indicating which surfaces should be sampled on which days.
  - (b) Surfaces to be tested must be cleaned and disinfected, dry, flat, sufficiently large and smooth.
  - (c) Three samples should be taken from large objects. Places that should receive most attention are the areas that may come into contact with the product. Approximately two thirds of the total number of samples should be taken from food contact surfaces.
  - (d) The following points should, for example, be chosen as sampling sites: knives (junction of blade and handle), hollow blood draining knives, elastrators, bung bagging machines, scraping/gambrelling table (pig), sawblades and cutters, cattle dehiding, other carcase dressing instruments, polishing machine, shackles and containers for transport, transport conveyor belts, aprons, cutting tables, flap doors, chutes for food organs, etc.

### Frequency

- 3. (a) A minimum of 10 samples (or up to 30 samples in a large production area) should be carried out within a period of two weeks.
  - (b) If the results are satisfactory over a period of time the frequency of sampling may be reduced following the agreement of the OVS, but fortnightly sampling must be resumed if unsatisfactory results are obtained.

# Sampling method

4. Either the Agar contact plate method or the swab technique may be used. In addition to the given descriptions, ISO methods may be used.

## (a) Agar contact plate method

- (i) Small plastic dishes with lids (i.e. internal diameter 5cm) filled with plate count agar (according to ISO, latest version) and dishes filled with violet red bile glucose agar (VRBG agar according to ISO, latest version) are pressed on to each sampling site and subsequently incubated. The contact surface of each plate is 20 cm<sup>2</sup>.
- (ii) Shortly before preparation of the plates, the relevant agar has to be melted to 100°C and cooled to 46 to 48°C. The plates have to be placed in a laminar air flow cabin and should be filled with agar until a convex surface is obtained. The prepared plates should be dried before use by incubating them upside down overnight at 37°C. This is also a useful check for possible contamination during preparation; plates with visible colonies must be discarded. After preparation the agar has a shelf life of approximately three months when kept at 2 to 4°C in closed bottles.
- (iii) The used contact plates do not need to be cooled during transport and before incubation. The plates have a shelf life of one week at 2 to 4°C, when sealed into plastic bags.

## (b) Swab technique

- (i) Samples should be collected with cotton swabs moistened with 1 ml of 0.1% NaCI peptone solution (8.5 g NaCI, 1g trypton casein-pepton, 0.1% agar, and 100 ml distilled water) from a surface area of preferably 20 cm<sup>2</sup>.
- (ii) If sampling is performed following cleaning and disinfection, an amount of 30g/litre Tween 80 and 3g/litre Lecithin (or other products with a similar effect) should be added to the moistening solution for swabs.
- (iii) The sampled surface must be swabbed 10 times from top to bottom applying a firm pressure on the surface.
- (iv) Swabs should be collected in a bottle containing 40 ml buffered peptone with 0.1% agar saline solution, then cooled and stored at 4°C until further processing.

(v) The bottle should be shaken vigorously before diluting in 10-fold steps in 40 ml 0.1% NaCI peptone solution followed by microbiological examination (e.g. drop-plating technique).

# Method for the examination of samples

- 5. (a) Analysis must be performed for total viable counts (TVC). Inoculated plate count agar plates and agar contact plates must be incubated for 24 hours at 37°C ± 1°C under aerobic conditions. This procedure must take place within two hours of sampling, unless this is impracticable, in which case it must take place as soon as is practicable after the end of that two-hour period. The number of bacterial colonies should be counted and recorded.
  - (b) Analysis for Enterobacteriaceae is voluntary unless required by the OVS. For quantitative estimation of Enterobacteriaceae VRBG agar must be used. Incubation of inoculated plates and agar contact plates must begin within two hours of sampling, unless this is impracticable, in which case it must take place as soon as is practicable after the end of that two-hour period. After 24 hours incubation at  $37^{\circ}\text{C} \pm 1^{\circ}\text{C}$  under aerobic conditions, the plates must be examined for Enterobacteriaceae growth.

#### **Records and results**

- 6. (a) The bacterial counts must be reported according to the number of organisms per cm<sup>2</sup> of surface area.
  - (b) Records must include:
    - (i) identification of the sample, date and time of sampling, name of the person that performed the sampling,
    - (ii) name and address of the laboratory which analysed the sample, date of investigation of samples in the laboratory, details of the method used and results.
  - (c) A responsible person from the laboratory should sign the records.
  - (d) Results have to be entered on a registration form and allocated into one of two categories established for the purpose of process control verification of cleaning and disinfection: "acceptable" and "unacceptable". The acceptable range for the number of colonies of TVC or Enterobacteriaceae are shown in the table.

# Values for the number of colonies for testing of surfaces

	Acceptable range	Unacceptable range
<b>Total viable counts (TVC)</b>	0 - 10/ cm <sup>2</sup>	>10/ cm <sup>2</sup>
Enterobacteriaceae	0 - 1/ cm <sup>2</sup>	>1/ cm <sup>2</sup>

# Feedback to staff

- 7. (a) The results of the test have to be reported to the responsible staff as soon as possible.
  - (b) The results should be used to maintain and improve the standard of cleaning and disinfection. Causes of unsatisfactory results should be clarified by consultation with the cleaning staff. The following factors may be involved: absence or inadequacy of training and/or instructions, the use of unsuitable cleaning and/or disinfection materials and chemicals, inadequate maintenance of cleaning apparatus, and inadequate supervision.".

# Amendments to the Poultry Meat, Farmed Game Bird Meat and Rabbit Meat (Hygiene and Inspection) Regulations (Northern Ireland) 1995

- **4.**—(1) The Poultry Meat Regulations are amended in accordance with the following paragraphs.
- (2) The following sub-paragraph is inserted between sub-paragraphs (aA) and (b) of paragraph (1) of regulation 8 (supervision of premises)—

- "(aB) the inspection of any documents and records required to be retained by the occupier pursuant to regulation 17(1)(f);".
- (3) In sub-paragraph (d) of paragraph (1) of regulation 17 (duties of occupier)—
  - (a) for "carry out" there shall be substituted "conduct regular"; and
  - (b) the word "and" at the end is revoked.
- (4) At the end of paragraph (1)(e)(iv) of regulation 17 there shall be added—
  "and
  - (f) retain for a period of at least 2 years any documents and records established by him in accordance with paragraph (4)(g).".
- (5) The following paragraph is added after paragraph (3) of regulation 17—
- "(4) The occupier of any licensed slaughterhouse used for slaughtering poultry, licensed cutting premises used for cutting up fresh poultry meat, licensed cold store used for the storage of fresh poultry meat or licensed re-wrapping centre used for packing, wrapping or re-wrapping fresh poultry meat shall conduct the regular checks on the general hygiene of conditions of production in those premises which are required by paragraph (1)(d) by implementing and maintaining a permanent procedure developed in accordance with the following principles—
  - (a) identify any hazards that must be prevented, eliminated or reduced to acceptable levels;
  - (b) identify the critical control points at the step or steps at which control is essential to prevent or eliminate a hazard or reduce it to acceptable levels;
  - (c) establish critical limits at critical control points which separate acceptability from unacceptability for the prevention, elimination or reduction of identified hazards;
  - (d) establish and implement effective monitoring procedures at critical control points;
  - (e) establish corrective actions when monitoring indicates that a critical control point is not under control;
  - (f) establish procedures to verify whether the measures outlined in sub-paragraphs (a) to (e) are working effectively; verification procedures shall be carried out regularly; and
  - (g) establish documents and records commensurate to the nature and size of the business to demonstrate the effective application of the measures outlined in sub-paragraphs (a) to (f) and to facilitate official controls."

Health, Social Services and Public Safety

Sealed with the Official Seal of the Department of Health, Social Services and Public Safety on 10th June 2002.

(L.S.) Denis McMahon

A Senior Officer of the Department of

#### **EXPLANATORY NOTE**

(This note is not part of the Regulations.)

These Regulations amend the Fresh Meat (Hygiene and Inspection) Regulations (Northern Ireland) 1997 (S.R. 1997 No. 493, as already amended) (*regulation 3*), and the Poultry Meat, Farmed Game Bird Meat and Rabbit Meat (Hygiene and Inspection) Regulations (Northern Ireland) 1995 (S.R.1995 No. 396, as already amended) (*regulation 4*).

These Regulations give effect to Commission Decision 2001/471/EC laying down rules for the regular checks on the general hygiene carried out by the operators in establishments according to Directive 64/433/EEC on health conditions for the production and placing on the market of fresh meat and Directive 71/118/EEC on health problems affecting the production and placing on the market of fresh poultry meat (O.J. No. L165, 21.6.2001, p. 48). They come into operation on 15th July 2002, except in relation to "small meat establishments" (defined in regulation 2(3)), in which case they come into operation on 7th June 2003.

The effect of the amendments made to the Fresh Meat (Hygiene and Inspection) Regulations (Northern Ireland) 1997 is that—

- (a) the occupier of any slaughterhouse, cutting premises, cold store or repackaging centre licensed under those Regulations is obliged to conduct the regular checks on the general hygiene of conditions of production in those premises which are already required by regulation 19(1)(d) of those Regulations by implementing and maintaining a permanent procedure developed in accordance with certain HACCP (Hazard Analysis and Critical Control Point) principles;
- (b) the occupier of a slaughterhouse licensed under those Regulations is obliged, in conducting the regular checks on the general hygiene of conditions of production in those premises which are referred to above, to carry out microbiological checks in relation to carcases and cleaning and disinfection of the premises; and
- (c) the occupier of cutting premises licensed under those Regulations is obliged, in conducting the regular checks on the general hygiene of conditions of production in those premises which are referred to above, to carry out microbiological checks in relation to cleaning and disinfection of the premises.

The effect of the amendments made to the Poultry Meat, Farmed Game Bird Meat and Rabbit Meat (Hygiene and Inspection) Regulations (Northern Ireland) 1995 is that the occupier of any slaughterhouse used for slaughtering poultry, cutting premises used for cutting up fresh poultry meat, cold store used for the storage of fresh poultry meat or rewrapping centre used for packing, wrapping or re-wrapping fresh poultry meat (in each case licensed under those Regulations) is obliged to conduct the regular checks on the general hygiene of conditions of production in those premises which are already required by regulation 17(1)(d) of those Regulations by implementing and maintaining a permanent procedure developed in accordance with certain HACCP principles.

£2.50

Published by The Stationery Office Limited

Printed in the UK by The
Stationery Office Limited
under the authority and
superintendence of Carol
Tullo, Controller of
Her Majesty's Stationery
Office being the Government
Printer for Northern Ireland and
the Officer appointed to print the
Acts of the Northern Ireland Assembly
Dd. N20530 C4. 6/02. Gp. 130. 14567.

