

SCHEDULE 2

TESTING METHODS

PART III

METHOD FOR THE ISOLATION OF *Enterobacteriaceae*

43. Tests shall be begun on receipt of the sample or on the first working day which allows this method to be completed. If the test is not begun on the day of receipt the sample shall be stored in a refrigerator until required at between 2°C and 8°C. If the sample has been refrigerated it shall be removed from the refrigerator and stored at room temperature for at least one hour before the test is started.

Samples

44. Tests shall be carried out using five 10 gram \pm 1 gram portions of each sample submitted for testing. Each 10 gram \pm 1 gram sample shall be placed aseptically in a sterile container containing 90 ml \pm 1 ml Buffered Peptone Water (BPW) and mixed thoroughly until the sample is evenly suspended.

Controls

45. Control tests shall be carried out each day that a test is initiated using –

- (a) *Escherichia coli* NCTC 10418 no more than 28 days old at time of use; and
- (b) processed animal protein or compost or digestive residue which is free of *Enterobacteriaceae*.

46. A 10 gram \pm 1 gram portion of the rendered animal protein shall be placed aseptically in a sterile container containing 90 ml \pm 1 ml BPW and mixed thoroughly until the sample is evenly suspended.

47. One colony of *Escherichia coli* shall be placed in 10 ml \pm 1 ml BPW and mixed to form an even suspension. Approximately 0.1 ml of the suspension shall be added to the suspension in the preceding paragraph.

48. This is then treated and examined in the same way as test samples. If no typical colonies are formed then that day's testing shall be invalid and shall be repeated.

Inoculations

49. For each portion of the sample 1 ml \pm 0.1 ml of solution shall be transferred to a sterile 90 mm petri dish (in duplicate). The plates shall be labelled to identify the portion of sample they were taken from. 15 ml \pm 1 ml of Violet Red Bile Glucose Agar (VRBGA)(1) at a temperature of 46°C \pm 1°C shall be added to each petri dish and immediately gently mixed by swirling the dish with five clockwise and five anticlockwise circular movements.

50. Once the agar has set, each agar plate shall be overlaid with a further 10 ml (approximately) of VRBGA at a temperature of 46°C \pm 1°C. Once the overlay has set, the plates shall be inverted and incubated aerobically at 37°C \pm 1°C for 20 hours \pm 2 hours.

(1) Violet Red Bile Glucose Agar – See Mossel, D.A.A., Eelderink, I. Koopmans, M., van Rossem, F. (1978) Laboratory Practice 27 No. 12 1049-1050; Emap Maclaren, PO Box 109, Maclaren House, 19 Scarbrook Road, Croydon CR9 1QH.

Status: This is the original version (as it was originally made). This item of legislation is currently only available in its original format.

Samples with colonies of *Enterobacteriaceae*

51. After incubation each set of duplicate plates shall be examined for colonies characteristic of *Enterobacteriaceae* (purple colonies 1-2 mm in diameter). All characteristic colonies on each plate shall be counted and the arithmetic mean of the duplicate plates taken.

The sample provisionally fails if either –

- (a) any arithmetic mean is above 30⁽²⁾; or
- (b) three or more arithmetic means are above 10;

in which case the following procedure shall be followed to establish whether or not the colonies are *Enterobacteriaceae*.

52. After counting the colonies, characteristic colonies shall be taken at random from the agar plates, the number being at least the square root of the colonies counted. Each of the colonies shall be subcultured onto a nutrient agar plate and incubated aerobically at $37^{\circ}\text{C} \pm 1^{\circ}\text{C}$ for 20 hours \pm 2 hours.

Examination of subcultures

53. An oxidase test and a glucose fermentation test shall be performed on each of the five subcultured colonies. Colonies which are oxidase-negative and glucose fermentation-positive shall be considered to be *Enterobacteriaceae*.

54. If not all of the colonies prove to be *Enterobacteriaceae*, the total count in paragraph 51 shall be reduced in proportion prior to establishing whether or not the sample should fail.

(2) An arithmetic mean of 30 is equivalent to 3×10^2 colony forming units per gramme of original sample